

EDITORIALS

- 1 I. Patient-centred outcomes in clinical research: does it really matter?
T. Heidegger, M. Nuebling, D. Saal and G. Kreienbühl
- 3 II. Thoracic epidural anaesthesia for cardiac surgery: are we missing the point?
T. M. Hemmerling, F. Carli and N. Noiseux
- 5 III. Volume 100
C. S. Reilly



REVIEW ARTICLE

- 8 Sedation and regional anaesthesia in the adult patient
D. Höhener, S. Blumenthal and A. Borgeat

SPECIAL ARTICLE

- 17 Anaesthesia chapter from *Saving Mothers' Lives; reviewing maternal deaths to make pregnancy safer*
G. M. Cooper and J. H. McClure



CARDIOVASCULAR

- 23 Meta-analysis of the effect of heart rate achieved by perioperative beta-adrenergic blockade on cardiovascular outcomes
B. M. Biccard, J. W. Sear and P. Foëx
- 29 Sevoflurane but not propofol increases interstitial glycolysis metabolites availability during tourniquet-induced ischaemia-reperfusion
M. Carles, J. Dellamonica, J. Roux, D. Lena, J. Levraut, J. F. Pittet, P. Boileau and M. Raucoules-Aime

CLINICAL PRACTICE

- 36 Pre-incisional epidural ropivacaine, sufentanil, clonidine, and (S)+ketamine does not provide pre-emptive analgesia in patients undergoing major pancreatic surgery
A. Gottschalk, M. Freitag, E. Steinacker, S. Kreißl, C. Rempf, H.-J. Staude, T. Strate and T. Standl
- 42 National survey of College Tutors in the UK regarding training in medical education
A. Rashid, A. Doger and G. Gould
- 45 Circadian distribution of sleep phases after major abdominal surgery
I. Gögenur, G. Wildschütz and J. Rosenberg
- 50 Comparison of surgical conditions during propofol or sevoflurane anaesthesia for endoscopic sinus surgery
H. J. Ahn, S.-K. Chung, H.-J. Dhong, H. Y. Kim, J. H. Ahn, S. M. Lee, T. S. Hahm and J. K. Kim

CRITICAL CARE

- 55 Effects of sodium nitroprusside on splanchnic microcirculation in a resuscitated porcine model of septic shock
A. Assadi, O. Desebbe, C. Kaminski, T. Rimmelé, F. Bénatir, J. Goudable, D. Chassard and B. Allaouchiche

NEUROSCIENCES AND NEUROANAESTHESIA

- 66 Regional cerebral metabolic rate (positron emission tomography) during inhalation of nitrous oxide 50% in humans
P. Reinstrup, E. Ryding, T. Ohlsson, A. Sandell, K. Erlandsson, K. Ljunggren, L. G. Salford, S. Strand and T. Uski
- 72 Effect-site half-time for burst suppression is longer than for hypnosis during anaesthesia with sevoflurane
R. R. Kennedy, C. Minto and A. Seethepalli

OBSTETRICS

- 78 Transcutaneous electrical nerve stimulation at the PC-5 and PC-6 acupoints reduced the severity of hypotension after spinal anaesthesia in patients undergoing Caesarean section
Y. C. P. Arai, N. Kato, M. Matsura, H. Ito, N. Kandatsu, S. Kurokawa, M. Mizutani, Y. Shibata and T. Komatsu

PAEDIATRICS

- 82 Variation of bispectral index under TIVA with propofol in a paediatric population
O. Tirel, E. Wodey, R. Harris, J. Y. Bansard, C. Ecoffey and L. Senhadji
- 88 Electrical velocimetry for measuring cardiac output in children with congenital heart disease
K. Norozi, C. Beck, W. A. Osthaus, I. Wille, A. Wessel and H. Bertram

PAIN

- 95 Efficacy of intravenous acetaminophen and lidocaine on propofol injection pain
O. Cambay, N. Celebi, O. Arun, A. H. Karagöz, F. Sarıcaoğlu and S. Özgen

REGIONAL ANAESTHESIA

- 99 Randomized controlled trial of patient-controlled epidural analgesia after orthopaedic surgery with sufentanil and ropivacaine 0.165% or levobupivacaine 0.125%
I. Smet, E. Vlamincx and M. Vercauteren
- 104 Spinal anaesthesia with articaine 5% vs bupivacaine 0.5% for day-case lower limb surgery: a double-blind randomized clinical trial
T. Dijkstra, J. A. Reesink, B. C. Verdouw, W. S. C. J. M. Van der Pol, T. Feberwee and A. G. Vulto
- 109 Epidural anaesthetic effect of the 8% emulsified isoflurane: a study in rabbits
Y.-F. Chai, J. Yang, J. Liu, H.-B. Song, J.-W. Yang, S.-L. Liu, W.-S. Zhang and Q.-W. Wang

RESPIRATION AND THE AIRWAY

- 116 Use of the McGrath® videolaryngoscope in the management of difficult and failed tracheal intubation
B. Shippey, D. Ray and D. McKeown
- 120 Upper cervical spine movement during intubation: fluoroscopic comparison of the AirWay Scope, McCoy laryngoscope, and Macintosh laryngoscope
K. Maruyama, T. Yamada, R. Kawakami, T. Kamata, M. Yokochi and K. Hara
- 125 Remifentanil target-controlled infusion vs propofol target-controlled infusion for conscious sedation for awake fiberoptic intubation: a double-blinded randomized controlled trial
M. R. Rai, T. M. Parry, A. Dombrowskis and O. J. Warner
- 131 Effects of carbon dioxide absorbent cooling and temperature gradient reduction on water condensation in the anaesthesia circuit
G. Hirabayashi, H. Uchino, T. Joko, H. Kaneko and N. Ishii
- 137 CORRESPONDENCE
- 144 BOOK REVIEWS
- 146 ERRATUM
- 147 ACKNOWLEDGEMENT OF ASSESSORS